




Regional climateprediction.net for the western US


Philip Mote
Oregon Climate Change Research Institute





climateprediction.net





Climate Experiment

board keys
e view


+ key when
over mode
ature
snow
e

art rotation
ide grid
more options

view:

surface

ter model of
n simulates
here & ocean
bal grid.
w shows the
scale. Switch
ud (C) and
ys to observe




Thanks for taking part! Modelling the first few years is extremely useful for

[the earth/climate change](#)


created by
climatechange

episodic
decades

Societal relevance
Scientific confidence




Spatial scale


An aerial photograph of a coastal town, likely in Japan, that has been severely flooded. The water is murky and brown, covering most of the land area. Several large industrial or commercial buildings are partially submerged. In the background, a large body of water, possibly a bay or harbor, is visible under a dramatic sky with dark, heavy clouds and bright sunbeams (crepuscular rays) breaking through. The overall scene conveys a sense of environmental disaster and the impact of extreme weather events.

Have heavy precipitation events increased?
Has the incidence of severe floods increased?
If so, is it due to global climate change?































projected increases in annual temperature



NARCCAP




NAARCCM

	GFDL	CGCM3	HADCM3	CCSM	NCEP
CRCM					
ECPC					
HRM3					
MM5I					
RCM3					
WRFP					


Winter Warming

1990s to 2050s
Temperature Change




Change in Winter Temperature (degrees C)

Difference between
regional and global models



Change in Winter Temperature (degrees C)





Year average of number of **frost days** per month

Number of days with **Tmax > threshold** (3 different) over the 10 years

Monthly **99th percentiles** of Tmax, **1st percentile** Tmin, computed over the 10 years

Maximum (Minimum) surface temperature per month and per year

Extreme precipitation

Maximum number of **consecutive dry days** over the 10 years

Average monthly **number of wet days** over the 10 years

Year **maximum 3-day precipitation**

Number of days with $P > \text{threshold}$ (3 different) over the 10 years

Monthly **99th high percentiles** of P, computed over the 10 years

Maximum daily P for each year

Extreme wind

Maximum daily wind speed for each year

Hydrological cycle

Year average of monthly soil moisture

Year average of 1st of each month **snow water equivalent**

Year average of monthly runoff

Yearly **maximum of daily runoff**

Maximum of maximum daily runoff in average over the 10 years

Time of last snow in average over the 10 years

Swelling

regional CPDN

- Higher spatial resolution
- Higher statistical resolution
- More societally relevant outputs